


C21.6

Antibodies


Structure & Function




Antibodies



- Immunoglobulin (Ig) is a gamma globulin protein that functions as an antibody. They are found in the blood plasma, tissue fluids, body secretions, on some leukocyte membranes but not inside our cytoplasm. Antibodies have receptors matched to foreign antigen.
- the basic structural unit of an antibody.....
- composed of four polypeptide chains linked by **disulfide (-S-S-) bonds**
- two larger **heavy chains** about 400 amino acids long // heavy chains have a hinge region where antibody is bent
- two **light chains** about half as long
- **variable (V) region** in all four chains // gives the antibody its uniqueness



Antibody Structure










Five Classes of Antibodies

- Remember “**MADGE**” (IgM – IgA – IgD – IgG – IgE)
 - Note: a single plasma cell is able to produce all four classes of antibodies – with same antigen binding site
 - Single plasma cells makes 2,000 antibodies per second for 7 days – clonal selection produces millions of active plasma cells
 - A single plasma cells may start to produce IgM antibodies and then produce IgG later in the infection




CLASS	FUNCTION
IgG 	<ul style="list-style-type: none">• Makes up the majority of antibodies in serum• The only antibody that can cross the placenta from mother to fetus• Functions in opsonization, neutralization, and complement fixation
IgA 	<ul style="list-style-type: none">• Found in secretions such as breast milk and saliva• Functions in agglutination and neutralization




CLASS	FUNCTION
IgM 	<ul style="list-style-type: none">• The first antibody secreted on exposure to an antigen• Potent agglutinating and precipitating agent• Functions in complement fixation
IgE 	<ul style="list-style-type: none">• Binds mast cells and basophils and triggers their degranulation, facilitating inflammation, particularly in the allergic response
IgD 	<ul style="list-style-type: none">• Antibody found exclusively on the surface of B cells• Has a role in B cell sensitization and activation


Agglutination



(a)



Antibody-dependent cell-mediated cytotoxicity (ADCC).




(a) Organisms, such as many parasites, that are too large for ingestion by phagocytic cells must be attacked externally.

FUNCTION

DESCRIPTION


Agglutination/precipitation




Antibodies clump antigens together to enhance phagocytosis.




Opsonization

IgG coats antigens and binds phagocytes, enhancing phagocytosis.



FUNCTION	DESCRIPTION
<p>Neutralization</p>  <p>The diagram shows several green Y-shaped antibody molecules. Each antibody has two orange triangular structures attached to its arms, which are labeled 'Toxin' with a line pointing to one of them. The antibodies are arranged in a way that their Y-shaped parts are facing the toxins, illustrating how they bind to and neutralize the pathogenic components.</p>	<p>Antibodies bind pathogenic components of toxins and block toxic effects.</p>
<p>Complement activation</p>  <p>The diagram illustrates the process of complement activation. It starts with a green Y-shaped antibody molecule. A blue oval labeled 'Complement protein' is shown binding to the antibody. An arrow points from this complex to a purple rod-shaped cell. On the surface of the cell, a cluster of blue dots is labeled 'Membrane attack complex (MAC)'. Another arrow points from the cell to a final stage where the cell is fragmented and surrounded by orange particles, labeled 'Lysis'.</p>	<p>Antibodies activate complement proteins, leading to cell lysis.</p>
<p>Stimulation of inflammation</p>  <p>The diagram shows a large, purple, irregularly shaped cell labeled 'Mast cell'. Several green Y-shaped molecules, labeled 'IgE', are attached to the surface of the cell. Small purple dots are shown inside the cell, and a cluster of these dots is labeled 'Inflammatory mediators'. An arrow points from the mediators inside the cell to a group of small purple dots being released from the cell, indicating the release of these mediators into the surrounding environment.</p>	<p>IgE binds mast cells and basophils, and triggers release of inflammatory mediators.</p>



Recognize

React

Remember

Born

Educated

Deploy

React

Remember

CELL-MEDIATED IMMUNITY
Directed against intracellular pathogens,
some cancer cells, and tissue transplants

ANTIBODY-MEDIATED IMMUNITY
Directed against extracellular pathogens

Humoral Immunity Responses



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.





Table 13.9 Primary and Secondary Response to Antigens

Upon the first exposure to an antigen, the system undergoes a **primary response**. The earliest part of this response, the *latent period*, is marked by a lack of antibodies for that antigen, but much activity is occurring. During this time, the antigen is being concentrated in lymphoid tissue and is being processed by the correct clones of B lymphocytes. As plasma cells synthesize antibodies, the serum titer increases to a certain plateau and then tapers off to a low level over a few weeks or months. Early in the primary response, most of the antibodies are the IgM type, which is the first class to be secreted by plasma cells. Later, the class of the antibodies (but not their specificity) is switched to IgG or some other class (IgA or IgE).

After the initial response, there is no activity, but memory cells of the same specificity are seeded throughout the lymphatic system.

When the immune system is exposed again to the same immunogen within weeks, months, or even years, a **secondary response** occurs. The rate of antibody synthesis, the peak titer, and the length of antibody persistence are greatly increased over the primary response. The speed and intensity seen in this response are attributable to the memory B cells that were formed during the primary response. The secondary response is also called the **anamnestic response**. The advantage of this response is evident: It provides a quick and potent strike against subsequent exposures to infectious agents.