

# Golgi tendon organ

From Wikipedia([View original Wikipedia article »](#)) Last modified on 22 November 2011 at 03:07


The **Golgi organ** (also called **Golgi tendon organ**, tendon organ, neurotendinous organ or neurotendinous spindle), is a **proprioceptive sensory receptor organ** that is located at the **insertion of skeletal muscle fibers into the tendons of skeletal muscle**. It provides the **sensory component of the Golgi tendon reflex**.

The Golgi organ should not be confused with the Golgi Apparatus, which is an organelle in the eukaryotic cell, or the Golgi stain, which is a histologic stain for neuron cell bodies.

## Anatomy

The body of the organ is made up of strands of collagen that are **connected at one end to the muscle fibers and at the other merge into the tendon proper**. Each tendon organ is **innervated by a single afferent type Ib sensory fiber that branches and terminates as spiral endings around the collagen strands**. The Ib afferent axon is a large diameter, myelinated axon. Each neurotendinous spindle is enclosed in a fibrous capsule which contains a number of enlarged tendon fasciculi (intrafusal fasciculi). One or more nerve fibres perforate the side of the capsule and lose their medullary sheaths; the axis-cylinders subdivide and end between the tendon fibers in irregular disks or varicosities (see figure).

## Function



Mammalian tendon organ showing typical position in a muscle (left), neuronal connections in spinal cord (middle) and expanded schematic (right). The tendon organ is a **stretch receptor that signals the force developed by the muscle**. The sensory endings of the Ib afferent are entwined amongst the musculotendinous strands of 10 to 20 motor units. Animated version: [1]

**When the muscle generates force, the sensory terminals are compressed.** This stretching deforms the terminals of the Ib afferent axon, opening stretch-sensitive cation channels. As a result, the Ib axon is depolarized and fires nerve impulses that are propagated to the spinal cord. The **action potential frequency signals the force being developed by the 10 to 20 motor units within the muscle**. This is representative of whole muscle force<sup>[1]</sup>

The Ib sensory feedback **generates spinal reflexes and supraspinal responses which control muscle contraction**. Ib afferents synapses with interneurons within the spinal cord that also project to the brain cerebellum and cerebral cortex. One of the main spinal reflexes associated with Ib afferent activity is the **autogenic inhibition reflex**, which helps regulate the force of muscle contractions. Tendon organs signal muscle force through the entire physiological range, not only at high strain.<sup>[1][2]</sup>

During locomotion, Ib input excites rather than inhibits motoneurons of the receptor-bearing muscles and it also affects the timing of the transitions between the stance and swing phases of locomotion.<sup>[3]</sup> The switch to autogenetic excitation is a form of positive feedback.<sup>[4]</sup>

The ascending or afferent pathways to the cerebellum are the dorsal and ventral spinocerebellar tracts. They are involved in the cerebellar regulation of movement.

## History

Until 1967 it was believed that Golgi tendon organs had a high threshold, only becoming active at high muscle forces. Consequently it was thought that tendon organ input caused "weightlifting failure" through the clasp-knife reflex, which protected the muscle and tendons from excessive force. However, the underlying premise was shown to be incorrect by James Houk and Elwood Henneman in 1967.<sup>[5]</sup>